



Denise Bickel DVM

Adult Horses:

Again, no more than 2 combo vaccines (4 way, 5 way, etc) or 3 individual vaccines should be administered at one time due to the serious risk of vaccine reactions

We only recommend the Intranasal (IN) strangles vaccine. We highly recommend that this vaccine be administered by a veterinarian due to the risk of cross-contamination and infection to the horse.

Adult horses that are not traveling or showing and do not have frequent contact with new or young horses:

Annually:

Tetanus
Rabies
EEE, WEE, VEE

Annually in Spring

West Nile
Potomac Horse Fever

Annually in July:

2nd Potomac Horse Fever

Adult horses that are traveling, showing, or have frequent contact with new or young horses:

Annually:

Tetanus
Rabies
EEE, WEE, VEE
IN strangles**

Twice yearly:

Rhino/Flu *

Annually in Spring:

West Nile
Potomac Horse Fever

Annually, July:

Potomac Horse Fever

* Horse under heavy competition or who have lots of contact with horses less than 3 years old require Rhino and Flu boosters every 3 months

** Horses who have lots of contact with horses less than 5 years old should have an IN strangles twice a year

Broodmares:

Annually, 4-6 weeks prepartum:

Tetanus
EEE, WEE, VEE
Rabies
West Nile
Potomac Horse Fever
Influenza

Annually pre-breeding

Influenza
Potomac Horse Fever

During Pregnancy at 5, 7, 9 months pregnant

Rhinopneumonitis vaccine specifically for broodmares

Please note that all vaccines carry a risk of reaction. We highly recommend a physical exam prior to yearly vaccinations to ensure the horse's health.